



Scleroderma and Pregnancy

You can get more information about Scleroderma from:

The Scleroderma Society

PO Box 581, Chichester, PO19 9EW Tel: 020 7000 1925

Published by The Scleroderma Society © 2008

Charity Registration Number 286736

www.sclerodermasociety.co.uk

[email:info@sclerodermasociety.co.uk](mailto:info@sclerodermasociety.co.uk)

Advice Line: 0800 311 2756

This leaflet is supported by a donation from
Actelion Pharmaceuticals UK Ltd

www.sclerodermasociety.co.uk

PO Box 581, Chichester, PO19 9EW Tel: 020 7000 1925

[email:info@sclerodermasociety.co.uk](mailto:info@sclerodermasociety.co.uk)

Advice Line: 0800 311 2756

Introduction:

The inter-relations between scleroderma and pregnancy are of considerable interest, particularly because scleroderma is a disease which affects markedly more females than males and does seem to have a peak of onset during the reproductive period. There are three questions of interest to those researching in the field. These are:

- Do women who are going to develop scleroderma have any difference in their pregnancy history prior to the development of the disease?
- Does the presence of scleroderma affect the outcome of pregnancy in any way?
- Does having a pregnancy affect the course or the outcome of scleroderma and its severity?

Pregnancy experience before the onset of Scleroderma:

There does seem to be a number of studies suggesting that in women who are going to develop scleroderma, there is a small increase in the proportion of these women reporting previous difficulty in conceiving. In some women this difficulty is such that they have no recorded successful conception prior to the disease. Further, information also suggests that the size of babies born to women who are going to develop scleroderma are

actually smaller than might be expected. It is difficult to know what these findings all mean, although there is some suggestion that Raynaud's Phenomenon, which is part of the scleroderma process, causes in some women a small reduction in blood flow to the uterus which can have these effects.

Recent studies have suggested that during a normal pregnancy cells from the baby can cross the placenta and circulate in the mother's bloodstream. In women who develop scleroderma, the baby's cells survive for longer and the reason for the development of the disease may be a response to these cells. Thus in the same way that a person who receives a transplanted kidney may react and become ill following the transplant, then the mother's scleroderma may be a very long delayed reaction to the persistence of these "foreign" cells.

What is the pregnancy experience after the onset of disease?

It used to be said that women who had developed scleroderma and then became pregnant, normally had a very poor experience with their pregnancy with a high rate of miscarriage and other pregnancy problems. However, more recent and more carefully done studies have actually shown that in women who have scleroderma and who become pregnant, the pregnancy loss rates are really only slightly higher than expected from normal

population experience. Indeed, the best study would suggest that there is no increase in miscarriage rates between women with scleroderma and the normal population women. As with pregnancies occurring before the onset of the disease, pregnancies after the onset of the disease are also associated with the risk of prematurity and slightly smaller babies. This does not appear to be a major problem in regard to the babies health, but it is of interest to those investigating the disease.

The Effect of Pregnancy on Scleroderma:

It is often recommended that women who had scleroderma should not become pregnant because this might cause deterioration in their disease. One of the problems for medical researchers is that when people have a bad experience, it tends to get published in the medical press, where in reality, one needs to answer whether this might have happened even if the women had not become pregnant. When carefully observed groups of women with scleroderma have been followed up and compared with what happened when they were not pregnant, the picture is much happier for the milder disease. The conclusions would be that for patients with limited scleroderma, there is no effect of pregnancy on the disease. However, women with diffuse scleroderma in pregnancy may have

problems with high blood pressure and some develop kidney disease. Further, some patients with scleroderma and lung disease deteriorate after delivery. It is therefore wise for each patient contemplating pregnancy to discuss their case individually with their doctor, so that the best advice can be obtained.

Conclusion:

There is a lot of hearsay and very little hard information on the relationship between pregnancy and scleroderma. There are some intriguing suggestions that pregnancy experience prior to disease onset might provide hints to understanding why women have an increased risk of this disorder. There is no evidence that reproductive loss through miscarriage and related factors is importantly increased either before or after the development of the disease. There are some suggestions however, that there is a slightly increased risk of prematurity, and that the birth weight of babies born from mothers with scleroderma might be lower. In some women who have scleroderma, pregnancy does not cause a problem although it is important carefully to monitor blood pressure, function of the kidneys and lung function during pregnancy and there is a need to attend regularly for specialist appraisal during this period.